

INHOUDSOPGAWE/CONTENTS

INHOUDSOPGAWE/CONTENTS.....	i
LYS VAN TABELLE/ LIST OF TABLES.....	ii
BELANGRIK	iv
DANKBETUIGINGS	v
IMPORTANT.....	vi
ACKNOWLEDGEMENTS	vii
KULTIVAREVALUERING.....	viii
KULTIVARINLIGTING	viii
DIAGNOSTIESE PARAMETERS.....	ix
KULTIVARS VOLGENS MIELIE PRODUKSIESTREKE	x
VRYWARING	x
CULTIVAR EVALUATION.....	xi
CULTIVAR INFORMATION.....	xi
DIAGNOSTIC PARAMETERS:	xii
CULTIVAR GROUPING ACCORDING TO MAIZE PRODUCTION REGIONS	xiii
INDEMNITY	xiv

LYS VAN TABELLE/ LIST OF TABLES

Tabel 1: Diagnostiese parameters vir die statistiese aanvaarbaarheid van proewe vir betroubare opbrengsanalises, 2015/2016 seisoen.....	1
Table 1: Diagnostic parameters for the statistical acceptability of trials for reliable yield analysis for the 2015/2016 season.....	1
Tabel 2: Gemiddelde graanopbrengs (t ha ⁻¹) vir mielie genotipes by verskillende omgewings in die oostelike gebiede gedurende die 2015/2016 seisoen	1,2,3
Table 2: Mean yield (t ha ⁻¹) for maize genotypes under different environments in the eastern region during the 2015/2016 season.....	1,2,3
Tabel 3: Gemiddelde graanopbrengs (t ha ⁻¹) vir mieliegenotipes by verskillende omgewings in die oostelike gebiede gedurende die 2014/2015 & 2015/2016 seisoene.....	4
Table 3: Mean yield (t ha ⁻¹) for maize genotypes under different environments in the eastern region during the 2014/2015 & 2015/2016 seasons	4
Tabel 4: Gemiddelde graanopbrengs (t ha ⁻¹) vir mieliegenotipes by oostelike omgewings gedurende die 2013/2014 & 2014/2015 & 2015/2016 seisoene	5,6,7
Table 4: Mean grain yield (t ha ⁻¹) for different maize genotypes at eastern environments during the 2013/2014 & 2014/2015 & 2015/2016 seasons	5,6,7
Tabel 5: Gemiddelde persentasie plante omval 2015/2016 seisoen.....	8
Table 5: Mean percentage lodged plants for the 2015/2016 season	8
Tabel 6: Gemiddelde persentasie spruitvorming 2015/2016 seisoen	9
Table 6: Mean percentage tillering for the 2015/2016 season	9
Tabel 7: Gemiddelde aantal koppe per plant 2015/2016 seisoen	10
Table 7: Mean number of ears per plant for the 2015/2016 season.....	10
Tabel 8: Gemiddelde graanvogpersentasie met oestyd 2015/2016 seisoen.....	11
Table 8: Mean percentage grain moisture at harvest for the 2015/2016 season	11
Tabel 9: Opsomming van gemiddelde waardes vir alle agronomiese parameters vir die 2015/2016 seisoen.....	12
Table 9: Summary of mean values for all agronomic parameters for the 2015/2016 season	12
Tabel 10: Meerjarige gemiddeldes ten opsigte van persentasie plante omval vir die 2013/2014, 2014/2015 & 2015/2016 seisoene	13
Table 10: Multi seasonal means for total percentage lodged plants for the 2013/2014,	

2014/2015 & 2015/2016 seasons	13
Tabel 11: Meerjarige gemiddeldes ten opsigte van persentasie spruitvorming vir die 2013/2014, 2014/2015 & 2015/2016 seisoene	14
Table 11: Multi seasonal means for total percentage tillering for the 2013/2014, 2014/2015 & 2015/2016 seasons	14
Tabel 12: Meerjarige gemiddeldes ten opsigte van aantal koppe per plant vir die 2013/2014, 2014/2015 & 2015/2016 seisoene	15
Table 12: Multi seasonal means for number of ears per plant for the 2013/2014, 2014/2015 & 2015/2016 seasons	15
Tabel 13: Meerjarige gemiddeldes ten opsigte van persentasie graanvog vir die 2013/2014, 2014/2015 & 2015/2016 seisoene	16
Table 13: Multi seasonal means for percentage grain moisture for the 2013/2014, 2014/2015 & 2015/2016 seasons	16
Tabel 14: Opsomming van gemiddelde waardes vir alle agronomiese parameters vir die 2013/2014, 2014/2015 & 2015/2016 seisoene.....	17
Table 14: Summary of mean values for all agronomic parameters for the 2013/2014, 2014/2015 & 2015/2016 seasons.....	17

BELANGRIK

Resultate van 'n kultivarproef by 'n enkele lokaliteit in enige jaar, of selfs 'n beperkte aantal lokaliteite in 'n enkele jaar, kan as gevolg van die kenmerkende seisoenale variasie in die Republic of South Africa (RSA) hoogs misleidend wees en kan sodoende onregverdiglik teen die beste genotypes vir daardie omgewing diskrimineer. **'N ERNSTIGE BEROEP WORD OP ALLE BETROKKNES GEDOEN OM NIE HUL GENOTIPEADVIES OP SO 'N HOOGS ONBETROUBARE METODE TE BASEER NIE.** Produsente word veral versoek om nougeset daarteen te waak dat hulle nie ook foutiewe genotipe uitsprake op dieselfde wyse doen nie, of op hierdie wyse mislei word nie.

Resultate van hierdie nasionale kultivarproewe, wat deur die LNR- Instituut vir Graangewasse uitgevoer is en gepubliseer word, geskied in belang van produsente, adviesdienste en die teeltbedryf. Die resultate mag derhalwe vryelik gebruik word, mits dit wetenskaplik korrek gedoen word deur die totale spektrum van lokaliteite en waarnemings in berekening te bring. Vrye gebruik van die resultate word ook met 'n verdere voorwaarde toegelaat, naamlik dat die nodige erkenning aan die bron van die inligting verleen word.

LNR-Instituut vir Graangewasse

Privaatsak X1251

POTCHEFSTROOM

2520

DANKBETUIGINGS

Die sukses van hierdie navorsingsprojek is toe te skryf aan die goeie samewerking en medewerking tussen die private en openbare sektore asook boere by wie genotipeproewe geplant is. Die verantwoordelike navorsers betuig hiermee hul grootste waardering vir die besondere samewerking en ondersteuning wat hulle van al die betrokkenes ontvang het.

Medewerkende Instansies

Medewerkers vir die 2015/2016 proefreeks word hier onder aangedui. **Hul getroue ondersteuning en uitstekende samewerking verdien vermelding en word erken.**

Agricol Saad (Edms) Bpk

Departement Landbou

Link Saad (Edms) Bpk

Klein Karoo

Monsanto SA (Edms) Bpk

Pannar Saad (Edms) Bpk

DuPont Pioneer RSA (Edms) Bpk

Seed-Co

LNR - Navorsingsinstituut vir Graangewasse

Hierdie verslag se samestelling, voorbereiding en vermeerdering het bydraes deur verskeie kollegas en beamptes geverg. Spesiale vermelding moet egter gemaak word aan Mnr. D De V Bruwer vir sy beplanning en bestuur van die proewe. Me T. Mathobisa-Manyokole vir data voorbereiding.

Die LNR-IGG wil hiermee ook sy dank uitspreek teenoor die Mielietrust vir hul finansiële ondersteuning wat die uitvoering van die proewe moontlik gemaak het.

IMPORTANT

Due to typical seasonal variations in the Republic of South Africa (RSA), results of a Cultivar trial at a single locality in any year, or even at a limited amount of localities in a single year can be highly misleading and can discriminate unfairly against genotypes which may in reality be the best for certain areas. **ALL THOSE INVOLVED, ARE STRONGLY URGED NOT TO BASE THEIR GENOTYPE RECOMMENDATIONS ON A HIGHLY UNRELIABLE METHOD SUCH AS THIS.** Producers, especially, are requested to avoid being misled in this way and against making incorrect genotype judgements.

The Agricultural Research Council (ARC) in the interest of producers, advisory services and the breeding industry publishes results of these national Cultivar trials, carried out by the ARC - Grain Crops Institute (GCI). These results may thus be freely used, as long as they are used in a scientifically correct manner, incorporating the whole spectrum of localities and observations. The source of the information should also be awarded the necessary recognition when using these results.

ARC-Grain Crops Institute

Private Bag X1251

POTCHEFSTROOM

2520

ACKNOWLEDGEMENTS

The success of this research project is a result of the good co-operation between the private, co-operative, and public sectors as well as farmers on whose farms cultivar trials were planted. The researchers wish to express their utmost appreciation for the exceptional co-operation and support received from all those involved.

Co-workers

The 2015/2016 trial series co-workers are listed below. **Their loyal support and excellent co-operation deserves mentioning and is acknowledged.**

Agricol Seed (Pty) Ltd

Department of Agriculture (DoA)

Link Seed (Pty) Ltd

Klein Karoo

Monsanto SA (Pty) Ltd

Pannar Seed (Pty) Ltd

DuPont Pioneer RSA (Pty) Ltd

Seed-Co

ARC - Grain Crops Institute

The compilation of this report was made possible by the efforts of many colleagues and officials. A special thanks goes to Mr D De V Bruwer for his planning and management of the trials. Ms. T. Mathobisa-Manyokole for data preparation

The ARC-GCI would like to thank the Maize Trust for their financial support that made these trials possible.

KULTIVAREVALUERING

Inligting rondom kultivars wat deur die produsent aangeplant word, is van kardinale belang. Dit beïnvloed 'n groot deel van die produsent se beplanning vir die seisoen. Betroubare en indien moontlik, onafhanklike inligting rakende kultivars moet aan elke produsent beskikbaar wees. Die LNR - Instituut vir Graangewasse (LNR-IGG), in samewerking met landboubesighede en die saadbedryf, poog om in hierdie belangrike behoefte van die produsent te voldoen. Die onus rus op die produsent om nuwe of onbekende kultivars eers op sy plaas te toets, voordat daar op groot skaal van beproefde kultivars afgesien word.

KULTIVARINLIGTING

Die boer moet aan die einde van elke seisoen 'n baie belangrike besluit neem naamlik: Watter kultivars gaan die volgende jaar geplant word? 'n Korrek beplande kultivarkeuse kan beslis 'n belangrike bydrae lewer om risiko te verminder en moet 'n belangrike onderdeel uitmaak van 'n produsent se produksiebeplanning.

Kultivars verskil van mekaar in een of meer van 'n verskeidenheid eienskappe en elkeen het dus 'n eiesoortige aanpasbaarheid en opbrengspotensiaal. Hierdie kultivar-verskeidenheid stel alternatiewes beskikbaar wat goed benut kan word en die volgende is belangrike riglyne vir kultivarbeplanning wat oorweeg moet word:

- Moet nooit 'n staatmaker-kultivar binne een seisoen met 'n nuwe of onbekende kultivar vervang nie.
- Kultivarverskeidenheid versprei risiko. Plant 'n reeks kultivars wat verskil in groeiseisoenlengte en eienskappe.
- Kultivars moet aanpas by 'n spesifieke opbrengs-potensiaal, maar moet in staat wees om hoër potensiaaltoestande te benut en 'n aanvaarbare opbrengs by laer potensiaaltoestande te lewer. Kultivars met 'n wye aanpassingsvermoë kan hier 'n rol speel.
- Hersien Kultivarkeuse jaarliks.

DIAGNOSTIESE PARAMETERS

- KV** Die Koëffisient van Variasie verwys na die fout van enkel persele en gee 'n aanduiding van die grootte van die variasie tussen perseelwaardes wat vanaf verskeie bronne afkomstig is. Die KV gee dus 'n aanduiding van die akkuraatheid van die perseelwaardes (grootte van die waarde). Bronne van variasie is byvoorbeeld grondvariasie (vrugbaarheid, diepte, grondvog, kleipersentasie, ongelykheid, ens) en plantvariasie (bevolkingsgrootte, oneweredige groei ens). Stremmingstoestande (vog, temperatuur, siektes, ens) het so dikwels tot gevolg dat normaalweg aanvaarbare grond- en plantvariasie baie sterker in die proefdata tot uiting kan kom en die KV vergroot. 'n Relatief hoë KV, wat aan hand van bekende bronne van variasie verklaar kan word, kan nie as die enigste parameter gebruik word om onbetroubare proefdata te identifiseer nie.
- GKV** Die Genetiese Koëffisient van Variasie verwys na die verskille in genotype-opbrengs. Die GKV is dus 'n aanduiding van die variasiegrootte wat aan verskille in genetiese samestelling tussen genotypeinskrywings toegeskryf kan word. Hoë waardes kan die gevolg wees van siektevatbaarheid, groot verskille in rypwordingstadium, temperatuurgevoeligheid en soortgelyke afwykings. Dit word ook gebruik om uitskieterproewe te identifiseer.
- tn** Genotipeherhaalbaarheid verwys na die herhaalbaarheid van genotypeGemiddeldes en kan gedefinieer word as die verwantskap tussen die genotype variasie en die totale variasie. Hierdie parameter is eintlik van waarde vir proewe waarvan die aantal herhalings nie dieselfde is nie.
- t** Die Intraklas Korrelasie verwys na die herhaalbaarheid van perseelwaardes oor herhalings. Hoe groter die ooreenstemming tussen perseelwaardes oor herhalings vir elke genotypeinskrywing is, hoe nader sal "t" na 1.0 neig.
- SF(t)** Die Standaard Fout van die Intraklas Korrelasie (t) gee 'n aanduiding van hoe akkuraat die skatting van "t" is.
- t/SF(t)**- Hierdie verhouding word as 'n belangrike parameter beskou daar die Intraklas Korrelasie (t) moet verkieslik minstens drie keer groter as sy foutterm moet wees. 'n Verhouding van kleiner as 3.0 dui aan dat die betrokke stel proefdata as minder betroubaar beskou kan word.

KULTIVARS VOLGENS MIELIE PRODUKSIESTREKE

Vir die doel van hierdie publikasie is die kultivarproewe ingedeel in streke volgens hitte eenhede, reënval en produksie en verder verwerk tot opbrengswaarskynlikheid. Waar voldoende inligting oor 'n betrokke kultivar beskikbaar is, is die inligting van twee en drie seisoene saamgevoeg. Andersins is die inligting van die nuwer kultivars bekom uit die eenjarige en tweejarige data van die onderskeie groter westelike streke. Om voldoende betroubare inligting te verskaf, is van die produksiegebiede saamgevoeg. Proewe wat in die oostelike produksiegebied gedoen is, is saamgevoeg (streek 2- warm oostelike gebied; streek 3- koue oostelike gebied en streek 4- KZN middelande).

Hierdie inligtingstuk poog nie om die enigste bron van inligting te wees nie. Kultivarinligting is ook by koöperasies en die saadmaatskappye beskikbaar.

VRYWARING

Die opsteller van die dokument en enige ander bron/instansie/persoon verantwoordelik vir enige inligting genoem in hierdie dokument is na die beste wete van die opstellers korrek met druktyd. Die inligting is ontwikkel deur wetenskaplike prosesse en word in goeder trou aangebied. Enige persoon/instansie wat hierdie inligting gebruik doen dit op eie risiko en die opstellers of enige ander party sal onder geen omstandighede verantwoordelik gehou kan word vir enige verliese gelei deur enige persoon/instansie wat die inligting in hierdie dokument gebruik nie.

CULTIVAR EVALUATION

Information on cultivars planted by producers is of utmost importance. It affects seasonal planning by producers to a large extent. Reliable and if possible independent information regarding cultivars should be available to every producer. The ARC- Grain Crops Institute (ARC-GCI) in co-operation with agricultural businesses and the seed industry attempt to satisfy the producers` requirements. The responsibility is with the producer to test new or unknown cultivars first before dispensing with the known cultivars on a large scale.

CULTIVAR INFORMATION

At the end of each season the farmer has to decide which cultivars are to be planted during the following season. A correctly planned cultivar choice can contribute greatly to reduce risk and constitutes an important part of the producer`s production planning.

Cultivars differ in one or more of a number of characteristics. Each cultivar has a particular adaptability and yield potential. Variability of cultivars provides alternatives, which can be utilized effectively. The following are important guidelines in cultivar planning which can be considered:

- Never replace a reliable cultivar with a new or unknown cultivar in a single season.
- Cultivar variability divides the risk factor. Use a series of cultivars, which differ in length of growing season and other characteristics.
- Cultivars must be adapted to a specific yield potential but should be able to utilize increased potential conditions and still deliver an acceptable yield at reduced yield potential. Cultivars with a wide adaptability can play a role under these conditions.
- Cultivar choice should be revised annually.

DIAGNOSTIC PARAMETERS:

CV The coefficient of variation - relates to the error of a single plot, and as such relates to the variability as induced by soil variation or plant population i.e. the larger the variation the larger the CV. Stress conditions (moisture, temperature, diseases, etc.) result in acceptable soil variation to be more pronounced in trials and a higher CV is recorded. The CV on its own cannot be used as a parameter to discard trials.

GCV The genetic coefficient of variation - relates to the yield differential between the highest and lowest entry yield, relative to the trial mean i.e. the greater the difference between the extreme values, the larger the GCV. High values are indicative of disease sensitivity, differences in maturity stage, temperature sensitivity and like problems.

tn Repeatability of genotype mean yield - relates to the repeatability of entry means, and can be defined as the relationship of genetic variance of observed means. In genotype trials this parameter is useful only when the number of replications between trials varies, otherwise the t-value is sufficient.

t The repeatability of plot yield or intra class correlation coefficient - relates to the repeatability of plot means over replications, and is interpreted as is the normal correlation coefficient, i.e. the greater the concurrence of plot values per entry over replications the closer "t" will strive towards unity. The standard error calculated for a particular t-value indicates the accuracy of the estimate of "t"

SE (t) Standard Error of the Intra-class Correlation (t) denotes how accurate the estimation of "t" is.

t/ SE (t) This relationship is considered an important parameter as the Inter Class Correlation (t) - Should be at least three times greater than its error term. A relationship of less the 3.0 denotes low reliability.

CULTIVAR GROUPING ACCORDING TO MAIZE PRODUCTION REGIONS

For the purpose of this publication the cultivar trials were divided into regions according to heat unit, precipitation and production for which yield reliability values were calculated. Information pertaining to three seasons was combined where sufficient information on a particular cultivar was available. Information on the newer cultivars was obtained from data of the annual and bi-annual reports of the greater eastern and western regions. Data for certain production regions have been combined in order to obtain more reliable information. Trials conducted in the eastern areas (region (1) warm eastern area; region (2) cold eastern area and region (4) KZN midlands) were combined.

Note that this brochure is not the only source of information. Cultivar information is also available at co-operatives and seed companies.

INDEMNITY

The composer of this document and any other source/institution/person responsible for any information contained in this document is to the best knowledge of the composers correct at printing. The information was developed using sound scientific procedures and is presented in good faith. Institutions or individuals can use this information at own risk and the composers or any other party will under no circumstances be under any legal obligation regarding any losses occurring by using the information contained in this document.

Tabel 1: Diagnostie parameters vir die statistiese aanvaarbaarheid van proewe vir betroubare opbrengsanalises vir die 2015/2016 seisoen

Table 1: Diagnostic parameters for the statistical acceptability of trials for reliable yield analysis for the 2015/2016 season

Omgewing Environment	Diagnostic parameters					
	SF SE	KV(%) CV%	GKV GCV	t t	SF(t) SE(t)	tn
Bapsfontein ⁽⁵⁾	0,62	16,60	6,30	0,13	0,09	0,31
Bethelhem ⁽¹⁾	0,56	18,30	9,50	0,21	0,09	0,44
Bloekomspruit ⁽¹⁾	0,40	12,20	15,40	0,61	0,07	0,82
Carolina ⁽⁷⁾	0,70	12,70	11,90	0,47	0,09	0,73
Cedara ⁽¹⁾	0,41	24,00	7,50	0,09	0,09	0,23
Delmas ⁽⁶⁾	0,60	13,30	14,80	0,55	0,08	0,79
Delmas ⁽³⁾	0,51	13,90	10,20	0,35	0,09	0,62
Dundee ⁽⁹⁾	0,92	24,10	14,50	0,26	0,09	0,51
Ermelo ⁽³⁾	0,41	21,30	3,60	0,03	0,09	0,08
Ermelo ⁽⁸⁾	0,37	17,60	30,80	0,75	0,05	0,90
Fouriesburg ⁽⁴⁾	0,50	14,90	12,00	0,40	0,09	0,67
Greytown ⁽⁸⁾	0,42	14,30	13,30	0,47	0,09	0,73
Kokstad ⁽⁹⁾	0,41	13,70	10,20	0,36	0,09	0,63
Middleburg ⁽³⁾	0,84	14,20	4,70	0,10	0,09	0,25
Petit ⁽³⁾	0,55	9,90	11,30	0,57	0,08	0,80
Wonderfontein ⁽²⁾	0,38	9,70	15,80	0,73	0,06	0,89

⁽¹⁾ARC; ⁽²⁾Pannar; ⁽³⁾Monsanto; ⁽⁴⁾Pioneer; ⁽⁵⁾Klein Karoo; ⁽⁶⁾DMS; ⁽⁷⁾Seedco; ⁽⁸⁾Link Seed; ⁽⁹⁾DoA

Tabel 2: Gemiddelde graanopbrengs (t ha⁻¹) vir mieliegenotipes by verskillende omgewings in die oostelike gebiede gedurende die 2015/2016 seisoen

Table 2: Mean yield (t ha⁻¹) for maize genotypes under different environments in the eastern region during the 2015/2016 season

Omgewing Environment	Genotipes /Genotypes									
	BG5285	DKC68-56R	DKC68-58BR	DKC71-42	DKC71-44B	DKC73-74BRGEN	DKC74-26R	DKC76-61B	DKC78-17B	DKC78-45BRGEN
Bapsfontein ⁽⁵⁾	7,20	6,25	7,28	7,16	7,92	7,71	6,36	6,52	6,00	7,62
Bloekomspruit ⁽¹⁾	6,74	5,46	6,54	4,21	5,54	6,41	6,96	5,68	6,07	7,27
Delmas ⁽³⁾	5,65	7,83	7,08	7,92	6,56	5,66	8,58	6,36	5,24	6,43
Delmas ⁽⁶⁾	8,80	8,35	7,46	8,38	7,09	7,21	5,00	8,75	7,79	6,12
Petit ⁽³⁾	10,20	10,53	10,05	10,19	9,68	10,24	11,22	8,53	9,43	8,52
Bethelhem ⁽¹⁾	6,78	5,33	6,28	4,42	5,79	4,61	5,69	5,97	5,72	5,87
Carolina ⁽⁷⁾	8,80	9,80	9,30	9,10	8,90	9,70	9,80	10,50	11,40	9,40
Ermelo ⁽⁸⁾	1,83	4,53	4,45	3,98	4,19	3,24	4,98	4,93	4,09	4,36
Ermelo ⁽³⁾	3,25	3,89	2,43	3,93	3,59	3,41	2,61	3,70	3,45	3,14
Fouriesburg ⁽⁴⁾	5,56	6,52	7,59	6,57	7,23	6,27	5,93	5,45	6,68	6,58
Middleburg ⁽³⁾	8,97	10,30	9,45	10,30	11,01	11,00	10,13	9,86	9,95	10,65
Wonderfontein ⁽²⁾	6,97	5,71	6,37	5,41	5,55	5,71	7,62	6,81	6,32	6,83
Cedara ⁽¹⁾	3,24	3,58	2,30	3,22	3,36	3,02	3,40	2,16	3,00	2,87
Greytown ⁽⁸⁾	4,71	5,67	4,77	5,52	6,66	6,14	6,26	5,59	6,55	6,17
Dundee ⁽⁹⁾	7,76	7,16	6,45	8,04	7,25	6,03	7,35	3,73	6,74	5,72
Kokstad ⁽⁹⁾	4,61	4,63	5,31	5,70	5,49	4,79	5,59	5,32	5,65	5,66
Gemiddelde/ Average (t ha ⁻¹)	6,32	6,60	6,44	6,50	6,61	6,32	6,72	6,24	6,50	6,45

⁽¹⁾ARC; ⁽²⁾Pannar; ⁽³⁾Monsanto; ⁽⁴⁾Pioneer; ⁽⁵⁾Klein Karoo; ⁽⁶⁾DMS; ⁽⁷⁾Seedco; ⁽⁸⁾Link Seed; ⁽⁹⁾DoA

Tabel 2 Vervolg.
Table 2 continue.

Omgewing Environment	Genotypes /Genotypes									
	DKC78- 79BR	DKC78- 87B	DKC80- 40BRG EN	IMP52- 11R	IMP52- 12R	IMP53- 49B	KKS830 1B	KKS832 6B	KKS840 8R	KKS841 OBR
Bapsfontein ⁽⁵⁾	6,34	7,20	8,56	5,86	5,74	6,73	6,54	5,81	5,55	6,08
Bloekomspruit ⁽¹⁾	7,61	6,80	7,08	6,15	5,28	5,57	4,01	3,99	5,80	5,90
Delmas ⁽³⁾	6,66	6,40	7,08	5,76	6,87	5,70	5,81	6,34	7,22	5,57
Delmas ⁽⁶⁾	7,30	8,83	7,27	6,26	6,36	7,76	6,53	5,96	7,36	9,58
Petit ⁽³⁾	11,43	9,89	11,37	11,63	8,33	9,61	8,58	9,31	9,61	9,81
Bethelhem ⁽¹⁾	5,85	6,31	5,67	4,96	4,12	6,11	3,83	5,55	5,07	5,28
Carolina ⁽⁷⁾	9,50	10,90	8,50	9,10	8,30	10,30	7,70	6,80	8,20	8,90
Ermelo ⁽⁸⁾	5,29	4,45	4,11	2,43	4,45	2,23	3,65	4,07	3,56	3,27
Ermelo ⁽³⁾	2,41	3,56	3,84	2,81	3,30	2,88	3,45	3,49	3,46	4,46
Fouriesburg ⁽⁴⁾	5,69	5,84	5,35	6,60	6,03	5,41	6,34	5,58	6,55	6,36
Middleburg ⁽³⁾	10,89	10,75	11,07	10,96	9,01	10,46	8,64	11,63	9,96	9,14
Wonderfontein ⁽²⁾	6,77	5,49	6,89	6,39	7,68	8,37	5,87	7,93	7,11	8,19
Cedara ⁽¹⁾	2,51	2,93	2,23	2,32	3,14	2,16	2,56	2,14	3,60	3,98
Greytown ⁽⁸⁾	5,04	5,26	5,94	4,72	4,68	5,72	5,69	4,69	5,80	5,15
Dundee ⁽⁹⁾	7,58	6,78	7,86	6,53	3,71	7,14	7,15	6,53	6,22	8,36
Kokstad ⁽⁹⁾	5,98	6,30	5,81	4,10	4,77	5,54	5,40	6,29	5,02	5,19
Gemiddelde/ Average (t ha ⁻¹)	6,68	6,73	6,79	6,04	5,74	6,36	5,73	6,01	6,26	6,58

⁽¹⁾ ARC; ⁽²⁾ Pannar; ⁽³⁾ Monsanto; ⁽⁴⁾ Pioneer; ⁽⁵⁾ Klein Karoo; ⁽⁶⁾ DMS; ⁽⁷⁾ Seedco; ⁽⁸⁾ Link Seed; ⁽⁹⁾ DoA

Tabel 2 Vervolg...
Table 2 continue...

Omgewing Environment	Genotypes/Genotypes									
	LS851 8	LS8526	LS8533R	LS8536B	LS8537	LS8539B	LS8541BR	P2137B	P2369WB	
Bapsfontein ⁽⁵⁾	5,41	5,62	6,76	5,89	6,78	6,84	5,10	6,97	6,61	
Bloekomspruit ⁽¹⁾	4,71	4,28	5,87	5,47	6,28	4,88	5,68	5,22	4,82	
Delmas ⁽³⁾	6,24	6,19	4,90	6,14	5,64	6,61	5,56	6,44	7,47	
Delmas ⁽⁶⁾	6,51	7,24	7,12	8,48	5,47	7,16	7,25	8,37	7,94	
Petit ⁽³⁾	8,32	9,00	7,85	8,68	9,02	7,97	9,55	12,59	11,05	
Bethelhem ⁽¹⁾	5,47	4,38	6,39	4,27	5,14	5,20	5,19	5,22	4,98	
Carolina ⁽⁷⁾	7,90	9,90	8,00	8,50	10,30	7,40	7,90	10,20	8,80	
Ermelo ⁽⁸⁾	3,87	1,84	3,06	3,96	2,29	2,84	3,23	5,56	4,01	
Ermelo ⁽³⁾	3,49	3,19	2,88	2,63	3,30	3,55	3,16	3,97	3,17	
Fouriesburg ⁽⁴⁾	6,52	5,03	4,97	6,86	5,58	5,18	4,12	6,08	6,22	
Middleburg ⁽³⁾	10,57	8,91	10,97	9,07	9,23	10,31	10,26	11,20	12,00	
Wonderfontein ⁽²⁾	4,27	9,01	5,05	6,07	9,53	6,63	6,68	5,88	5,62	
Cedara ⁽¹⁾	3,27	2,94	3,02	2,91	3,14	2,84	2,67	2,60	3,21	
Greytown ⁽⁸⁾	4,41	3,28	4,62	4,00	4,14	4,42	3,88	6,48	5,23	
Dundee ⁽⁹⁾	7,00	7,08	5,93	6,79	6,67	8,39	7,22	6,75	4,40	
Kokstad ⁽⁹⁾	4,48	5,26	4,11	4,11	4,31	5,05	5,18	6,04	5,76	
Gemiddelde / Average (t ha ⁻¹)	5,78	5,82	5,72	5,86	6,05	5,95	5,79	6,85	6,33	

⁽¹⁾ ARC; ⁽²⁾ Pannar; ⁽³⁾ Monsanto; ⁽⁴⁾ Pioneer; ⁽⁵⁾ Klein Karoo; ⁽⁶⁾ DMS; ⁽⁷⁾ Seedco; ⁽⁸⁾ Link Seed;
⁽⁹⁾ DoA

Tabel 2 Vervolg.
Table 2 continue....

Omgewing Environment	Genotipes/Genotypes										
	P243 2R	P2653 WB	PAN4A- 111	PAN4A- 172	PAN4B- 311B	PAN4R- 511R	PAN5A- 182	PAN5A- 291	PAN5R- 791BR	PAN6P- 110	PAN6Q -245
Bapsfontein ⁽⁵⁾	5,55	7,52	6,19	6,30	5,65	5,34	6,83	5,87	6,04	6,96	6,55
Bloekomspruit ⁽¹⁾	5,51	6,40	5,77	4,69	4,97	6,15	5,96	5,25	6,03	5,56	5,99
Delmas ⁽³⁾	6,62	6,53	5,17	6,48	5,73	7,17	7,13	5,97	5,69	5,39	5,87
Delmas ⁽⁶⁾	8,72	8,10	6,90	9,74	8,42	8,90	9,24	9,21	9,40	10,27	8,49
Petit ⁽³⁾	11,47	9,42	10,09	10,00	11,59	12,57	11,08	8,24	9,75	9,53	8,94
Bethelhem ⁽¹⁾	5,81	5,97	6,12	3,53	4,88	5,29	7,02	5,99	5,39	4,90	5,52
Carolina ⁽⁷⁾	8,50	11,00	8,70	8,10	7,50	8,70	11,80	11,90	9,80	11,10	9,70
Ermelo ⁽⁸⁾	4,38	2,63	2,00	6,12	2,76	2,82	4,94	3,16	2,36	6,88	2,24
Ermelo ⁽³⁾	3,12	3,71	3,04	3,75	3,10	3,45	3,94	3,18	3,57	3,62	3,73
Fouriesburg ⁽⁴⁾	6,06	6,58	7,03	6,55	5,32	6,19	7,19	5,73	6,24	3,97	4,83
Middleburg ⁽³⁾	10,36	10,79	11,20	12,15	9,75	11,99	10,16	10,64	11,79	9,93	11,10
Wonderfontein ⁽²⁾	6,38	9,04	6,84	6,59	7,20	6,94	6,61	6,90	6,65	8,20	7,91
Cedara ⁽¹⁾	3,40	2,94	2,49	3,32	3,38	3,21	2,20	2,80	2,41	3,17	2,90
Greytown ⁽⁸⁾	5,90	4,95	4,30	5,53	4,26	3,70	5,05	5,43	5,18	5,55	4,37
Dundee ⁽⁹⁾	7,73	6,62	6,99	2,06	5,84	7,85	7,61	8,43	9,07	6,92	7,28
Kokstad ⁽⁹⁾	5,18	4,45	4,17	6,31	5,48	4,23	5,48	4,72	4,79	5,74	4,71
Gemiddelde Average (t ha ⁻¹)	6,54	6,67	6,06	6,33	5,99	6,53	7,01	6,46	6,51	6,73	6,26

⁽¹⁾ ARC; ⁽²⁾ Pannar; ⁽³⁾ Monsanto; ⁽⁴⁾ Pioneer; ⁽⁵⁾ Klein Karoo; ⁽⁶⁾ DMS; ⁽⁷⁾ Seedco; ⁽⁸⁾ Link Seed; ⁽⁹⁾ DoA

Tabel 2 Vervolg.
Table 2 continue.

Omgewing Environment	Genotipes/Genotypes										Gemiddelde Average (t ha ⁻¹)
	PAN6Q- 345CB	PAN6Q -408CB	PAN6R -680R	PAN6R- 880CBGT	Phb31 M09	Phb33 H52B	SC402	SC419	SC506	SC512	
Bapsfontein ⁽⁵⁾	7,09	7,02	5,74	6,86	6,26	6,49	5,45	6,61	5,74	6,96	6,47
Bloekomspruit ⁽¹⁾	5,96	6,28	7,42	6,68	4,66	4,27	4,42	4,52	4,22	4,15	5,62
Delmas ⁽³⁾	5,92	7,62	6,07	6,34	7,42	7,87	5,00	5,68	6,58	5,81	6,36
Delmas ⁽⁶⁾	8,34	10,41	8,65	8,60	7,39	9,28	6,39	6,62	5,81	5,46	7,76
Petit ⁽³⁾	10,26	8,54	7,73	10,26	9,59	8,97	8,86	8,83	8,04	7,95	9,68
Bethelhem ⁽¹⁾	5,77	5,90	5,91	5,25	4,52	3,94	4,51	4,58	4,85	5,23	5,33
Carolina ⁽⁷⁾	10,80	10,90	12,60	9,30	8,70	9,10	7,50	7,50	9,40	8,80	9,30
Ermelo ⁽⁸⁾	3,35	3,93	3,36	5,15	2,99	4,11	2,31	1,63	2,61	2,34	3,62
Ermelo ⁽³⁾	3,34	3,48	3,31	3,55	3,47	3,04	2,53	3,15	4,08	3,50	3,36
Fouriesburg ⁽⁴⁾	4,07	4,61	4,59	6,54	7,05	6,21	4,97	4,52	5,16	5,80	5,88
Middleburg ⁽³⁾	10,32	9,21	9,66	7,60	10,57	10,83	9,29	8,66	9,81	9,87	10,25
Wonderfontein ⁽²⁾	7,20	7,45	7,64	8,92	6,36	6,31	5,33	8,64	7,35	5,15	6,85
Cedara ⁽¹⁾	3,28	3,01	3,28	3,64	2,38	2,89	2,58	3,92	3,00	2,46	2,94
Greytown ⁽⁸⁾	5,67	6,28	5,13	4,47	4,54	5,94	4,71	3,83	5,52	5,31	5,14
Dundee ⁽⁹⁾	5,39	7,74	5,63	7,68	6,17	6,48	5,75	6,66	5,66	3,99	6,64
Kokstad ⁽⁹⁾	5,26	5,22	5,34	5,71	4,24	6,64	4,66	4,04	5,72	4,67	5,16
Gemiddelde / Average (t ha ⁻¹)	6,38	6,72	6,38	6,66	6,02	6,40	5,27	5,59	5,85	5,47	

⁽¹⁾ ARC; ⁽²⁾ Pannar; ⁽³⁾ Monsanto; ⁽⁴⁾ Pioneer; ⁽⁵⁾ Klein Karoo; ⁽⁶⁾ DMS; ⁽⁷⁾ Seedco; ⁽⁸⁾ Link Seed; ⁽⁹⁾ DoA

Tabel 3: Gemiddelde graanopbrengs (t ha⁻¹) vir mieliegenotipes by verskillende omgewings in die oostelike gebiede gedurende die 2014/2015 & 2015/2016 seisoene

Table 3: Mean yield (t ha⁻¹) for maize genotypes under different environments in the eastern region during the 2014/2015 & 2015/2016 seasons

Omgewing/ Environment	Genotipes/ Genotypes										Gemiddelde/ Average (t ha ⁻¹)
	IMP 52-12R	KKS 8301B	KKS 8326B	LS8537	P2137B	PAN4A- 111	PAN4R- 511R	PAN5A- 182	PAN5R- 791BR	KKS 8301B	
2014/2015											
season											
Bapsfontein ⁽⁵⁾	4,99	6,31	3,75	4,93	5,02	5,07	4,26	6,16	4,84	5,47	5,08
Bergville ⁽¹⁰⁾	7,40	7,08	8,49	9,85	8,29	9,18	9,48	8,44	9,09	9,38	8,67
Bergville ⁽²⁾	11,76	12,33	13,66	12,57	14,17	13,46	13,44	12,90	13,45	11,4	12,92
Bethlehem ⁽¹⁾	5,25	4,39	5,50	6,37	4,78	5,35	6,06	5,07	6,03	5,67	5,45
Carolina ⁽⁷⁾	9,85	8,86	8,51	10,42	11,16	9,10	10,36	10,32	9,46	7,49	9,55
Delmas ⁽³⁾	10,79	7,64	7,60	10,11	8,45	9,58	10,49	10,31	8,26	9,79	9,30
Delmas ⁽²⁾	8,88	7,41	8,62	9,81	8,46	9,13	9,02	9,79	9,89	9,19	9,02
Devon ⁽²⁾	4,59	4,91	5,74	0,00	5,53	6,09	5,68	5,53	5,75	5,59	4,94
Greytown ⁽²⁾	4,75	5,16	5,59	4,82	6,12	1,61	5,52	5,71	5,53	5,59	5,04
Kinross ⁽⁷⁾	7,78	6,36	6,75	6,16	7,85	8,63	7,54	8,99	7,99	5,15	7,32
Middleburg ⁽³⁾	9,98	9,92	10,08	12,64	11,66	12,47	11,18	12,32	12,17	11,44	11,39
Moorivier ⁽⁷⁾	8,33	6,47	6,79	6,69	8,05	9,50	8,75	8,41	7,78	8,16	7,89
Standerton ⁽²⁾	3,66	3,76	4,08	3,79	3,85	4,19	4,37	4,36	4,00	3,58	3,96
Vrede ⁽⁷⁾	4,16	3,24	4,35	3,95	4,41	5,96	3,73	5,02	2,00	6,68	4,35
Winterton ⁽⁷⁾	12,74	12,14	13,47	13,51	13,72	12,38	14,68	12,14	12,37	12,31	12,95
Wonderfontein ⁽⁴⁾	12,34	10,46	10,10	13,03	13,41	13,27	12,51	14,54	12,25	13,4	12,53
2015/2016											
season											
Bapsfontein ⁽⁵⁾	5,74	6,54	5,81	6,78	6,97	6,19	5,34	6,83	6,04	6,73	6,30
Bethlehem ⁽¹⁾	4,13	3,83	5,55	5,13	5,22	6,13	5,29	7,02	5,39	6,11	5,38
Bloekomspruit ⁽¹⁾	5,28	4,01	3,99	6,28	5,22	5,77	6,15	5,96	6,03	5,57	5,43
Carolina ⁽⁷⁾	8,33	7,64	6,79	10,33	10,23	8,70	8,72	11,77	9,80	10,31	9,26
Cedara ⁽¹⁾	1,77	1,56	1,79	1,47	0,86	1,74	1,54	1,71	1,29	1,4	1,51
Delmas ⁽³⁾	6,36	6,52	5,96	5,47	8,37	6,90	8,90	9,24	9,40	7,7	7,48
Delmas ⁽³⁾	6,87	5,81	6,34	5,64	6,44	5,17	7,17	7,13	5,69	5,7	6,19
Dundee ⁽⁹⁾	3,71	7,15	6,53	6,67	6,75	6,99	7,85	7,61	9,07	7,14	6,95
Ermelo ⁽⁸⁾	4,46	3,65	4,07	2,29	5,56	3,00	2,82	4,94	2,36	2,23	3,54
Ermelo ⁽³⁾	3,30	3,45	2,44	3,30	3,97	3,04	3,46	3,94	3,57	2,88	3,34
Fouriesburg ⁽⁴⁾	6,02	6,34	5,58	5,58	7,82	7,03	6,19	7,19	6,24	5,41	6,34
Greytown ⁽⁸⁾	4,68	5,69	4,69	4,14	6,48	4,30	3,70	5,05	5,18	5,72	4,96
Kokstad ⁽⁹⁾	4,77	5,40	6,29	4,31	6,04	4,17	4,23	5,49	4,80	5,54	5,10
Middelburg ⁽³⁾	9,01	8,64	11,64	9,22	11,20	11,21	11,99	10,16	11,78	10,46	10,53
Petit ⁽³⁾	8,32	8,58	9,31	9,02	12,59	10,09	10,99	11,09	9,75	9,61	9,93
Wonderfontein ⁽²⁾	6,07	5,88	7,35	9,53	5,89	6,84	6,94	6,61	6,65	5,88	6,76
Gemiddelde/ Average (t ha ⁻¹)	6,75	6,47	6,79	6,99	7,64	7,26	7,45	7,87	7,31	6,47	

⁽¹⁾ ARC; ⁽²⁾ Pannar; ⁽³⁾ Monsanto; ⁽⁴⁾ Pioneer; ⁽⁵⁾ Klein Karoo; ⁽⁷⁾ Seedco; ⁽⁸⁾ Link Seed; ⁽⁹⁾ DoA; ⁽¹⁰⁾ Agricol

Tabel 4: Gemiddelde graanopbrengs (t ha⁻¹) vir mieliegenotipes by oostelike omgewings gedurende die 2013/2014, 2014/2015 & 2015/2016 seisoene

Table 4: Mean grain yield (t ha⁻¹) for different maize genotypes at eastern environments during the 2013/2014, 2014/2015 & 2015/2016 seasons

Omgewing/ Environment	Genotipes/Genotypes									
	BG5285	DKC73- 74BRGEN	DKC78- 17B	DKC78- 45BRGEN	DKC78- 79BR	DKC78- 87B	DKC80- 40BRGEN	IMP52- 11R	KKS8410 BR	LS8518
2013/2014 season										
Babsfontein ⁽⁵⁾	11,29	10,40	10,23	9,77	11,94	10,61	10,34	10,77	8,05	10,84
Bergville ⁽²⁾	6,48	5,93	6,01	6,96	6,26	6,14	5,59	6,70	6,32	5,53
Bethlehem ⁽¹⁾	13,25	10,57	14,32	11,65	13,17	13,68	11,74	11,97	10,42	11,02
Bethlehem ⁽³⁾	4,91	7,49	7,52	6,90	5,40	6,76	6,82	5,85	6,27	6,02
Bloekomspruit ⁽¹⁾	8,64	9,34	9,16	8,93	9,81	9,97	9,32	9,45	7,93	8,68
Carolina ⁽⁷⁾	6,42	7,52	7,32	6,78	7,58	7,46	7,28	6,63	8,33	6,97
Delmas ⁽²⁾	11,48	9,24	9,80	10,45	11,67	10,65	11,08	11,96	10,35	11,38
Delmas ⁽³⁾	13,09	9,97	13,11	10,91	12,31	11,84	11,85	11,86	10,72	12,11
Devon ⁽⁸⁾	5,73	6,22	6,81	7,10	6,66	7,11	6,48	4,95	6,22	4,97
Dundee ⁽⁹⁾	7,09	7,78	6,77	7,18	5,65	7,27	7,16	6,65	6,85	7,74
Ermelo ⁽⁴⁾	11,89	12,47	12,30	11,42	13,36	12,95	11,82	11,61	9,39	10,19
Ermelo ⁽¹⁰⁾	8,12	9,40	8,64	8,44	8,64	8,84	9,44	10,50	10,02	7,74
Greytown ⁽²⁾	10,69	10,54	11,12	10,14	10,30	10,02	10,03	10,19	9,86	10,72
Greytown ⁽²⁾	9,63	8,89	8,71	8,20	8,79	7,86	9,36	9,10	9,05	9,76
Hendrina ⁽³⁾	9,66	9,43	9,28	8,14	9,24	7,40	8,85	8,54	8,33	8,15
Kinross ⁽⁸⁾	11,73	11,08	11,41	10,52	10,64	10,78	10,94	10,41	9,80	9,04
Middelburg ⁽⁵⁾	10,44	9,13	8,48	10,53	9,29	7,57	7,76	9,12	8,09	8,79
Moorivier ⁽⁷⁾	9,60	9,46	10,16	11,13	9,29	9,54	12,93	11,37	12,13	10,21
Ogies ⁽³⁾	12,06	11,53	12,53	11,90	13,57	13,15	11,99	12,61	10,16	12,20
Petit ⁽³⁾	10,44	11,51	10,35	10,21	9,76	9,87	10,41	8,95	8,60	10,20
Villiers ⁽⁸⁾	11,41	10,06	12,14	9,62	9,49	10,10	10,01	10,27	8,03	9,87
Winterton ⁽⁷⁾	15,34	14,15	13,64	13,69	13,28	13,86	14,18	14,26	11,78	15,15
Wonderfontein ⁽²⁾	7,90	7,04	7,61	9,14	7,46	6,81	6,35	8,01	7,52	7,23
2014/2015 season										
Babsfontein ⁽⁵⁾	5,22	6,74	5,34	5,63	4,85	5,22	6,17	4,97	5,70	5,34
Bergville ⁽¹⁰⁾	8,09	9,32	8,72	6,66	8,00	9,11	10,48	8,33	8,21	9,93
Bergville ⁽²⁾	11,79	13,02	12,42	8,99	12,50	11,84	12,77	10,39	11,91	12,49
Bethlehem ⁽¹⁾	6,41	5,61	6,27	5,91	6,08	5,18	5,97	5,19	4,23	5,96
Carolina ⁽⁷⁾	11,40	10,36	9,88	8,96	10,05	10,88	9,43	9,48	10,48	9,35
Delmas ⁽³⁾	7,64	8,68	10,43	7,88	9,90	9,30	8,63	6,57	11,04	10,06
Delmas ⁽²⁾	8,49	9,53	8,88	8,92	8,74	10,82	8,95	9,43	9,55	9,43
Devon ⁽⁷⁾	6,72	6,09	5,29	5,07	6,80	4,14	4,93	6,06	5,99	5,87
Greytown ⁽²⁾	5,81	6,90	6,77	6,21	7,16	7,10	7,08	5,08	5,63	5,94
Kinross ⁽⁸⁾	8,18	7,93	5,38	7,60	7,06	7,86	8,83	8,12	8,21	7,17
Middelburg ⁽⁵⁾	12,84	11,08	11,24	11,64	11,58	11,54	12,36	7,87	10,51	11,07
Moorivier ⁽⁷⁾	7,97	7,41	8,64	8,44	6,40	6,26	10,16	8,53	8,57	8,30
Standerton ⁽²⁾	3,08	3,78	3,29	3,17	3,65	4,06	3,39	3,83	3,77	3,71
Vrede ⁽⁸⁾	3,71	3,42	5,83	0,00	4,50	4,50	6,00	5,81	3,78	5,72
Winterton ⁽⁷⁾	13,32	13,26	13,62	13,04	13,29	12,62	12,61	11,51	12,75	12,71
Wonderfontein ⁽²⁾	11,65	12,43	9,82	12,76	13,19	12,33	11,66	13,12	12,41	12,96
2015/2016 season										
Babsfontein ⁽⁵⁾	7,19	7,71	6,01	7,62	6,34	7,20	8,57	5,86	6,08	5,41
Bethlehem ⁽¹⁾	6,78	4,61	5,72	5,87	5,85	6,31	5,67	4,97	5,28	5,47
Bloekomspruit ⁽¹⁾	6,74	6,42	6,07	7,27	7,61	6,80	7,08	6,14	5,90	4,71
Carolina ⁽⁷⁾	8,84	9,74	11,44	9,43	9,44	10,94	8,55	9,10	8,88	7,88
Cedara ⁽¹⁾	3,24	3,02	3,00	2,88	2,51	2,93	2,23	2,32	3,98	3,28
Delmas ⁽³⁾	7,44	7,21	7,80	4,02	7,30	8,83	7,26	6,26	9,58	6,51
Delmas ⁽⁶⁾	5,65	5,66	5,25	6,43	6,66	6,40	7,08	5,77	5,57	6,24
Dundee ⁽⁹⁾	7,73	5,98	6,67	5,66	7,60	6,72	7,84	6,54	8,29	6,98
Ermelo ⁽⁸⁾	3,25	3,41	3,45	3,14	2,41	3,56	3,84	2,81	4,46	3,49
Ermelo ⁽³⁾	1,83	3,24	4,09	4,36	5,29	4,45	4,11	2,43	3,27	3,87
Fouriesburg ⁽⁴⁾	5,56	6,27	6,68	6,58	5,69	5,84	5,35	6,60	6,36	6,52
Greytown ⁽⁸⁾	4,71	6,14	6,55	6,17	5,04	5,26	5,94	4,73	5,14	4,41
Kokstad ⁽⁹⁾	4,61	4,79	5,65	5,66	5,98	6,31	5,81	4,10	5,19	4,48
Middelburg ⁽³⁾	8,97	10,99	9,96	7,05	10,90	10,75	11,07	10,95	9,14	10,57
Petit ⁽³⁾	10,20	10,24	9,43	8,52	11,43	9,90	11,37	11,63	9,81	8,32
Wonderfontein ⁽²⁾	6,97	7,64	6,32	6,83	6,77	5,49	6,89	6,39	8,19	4,27
Gemiddelde/ Average (t ha ⁻¹)	8,35	8,32	8,42	7,96	8,44	8,38	8,54	8,05	8,04	8,05

(¹) ARC; (²) Pannar; (³) Monsanto; (⁴) Pioneer; (⁵) Klein Karoo; (⁷) Seedco; (⁸) Link Seed; (⁹) DoA; (¹⁰) Agricol

Tabel 4 Verilog..
Table 4 Continue...

Omgenwing/ Environment	Genotypes/Genotypes									
	LS8526	LS 8533R	LS 8536B	LS 8539B	P2432R	P2653 WB	PAN4B- 311B	PAN6P- 110	PAN6Q- 245	PAN6Q- 345CB
2013/2014 season										
Babsfontein ⁽⁵⁾	9,29	8,34	9,83	9,71	11,45	10,23	9,97	8,66	11,61	12,36
Bergville ⁽²⁾	6,31	6,30	5,11	5,96	6,24	5,97	6,55	6,84	6,28	6,20
Bethlehem ⁽¹⁾	12,11	11,44	12,33	10,60	13,13	11,92	10,97	13,53	12,61	11,76
Bethlehem ⁽³⁾	7,07	6,21	7,72	4,45	6,66	6,23	7,19	5,00	5,80	5,05
Bloekomspruit ⁽¹⁾	8,21	8,68	8,41	8,31	9,13	9,27	10,17	9,25	9,38	10,02
Carolina ⁽⁷⁾	6,64	7,62	7,85	7,70	6,58	7,03	6,70	8,21	6,79	7,02
Delmas ⁽²⁾	10,25	10,71	10,16	9,98	11,75	10,57	11,77	10,28	9,73	9,61
Delmas ⁽³⁾	10,06	10,66	11,49	12,51	12,39	13,12	12,28	12,43	12,04	11,94
Devon ⁽⁸⁾	5,70	6,38	5,37	7,66	6,32	5,74	7,17	6,76	7,14	4,59
Dundee ⁽⁹⁾	6,14	7,10	6,73	5,45	5,93	7,28	7,21	6,93	6,73	6,62
Ermelo ⁽⁴⁾	10,27	10,69	10,02	10,50	11,13	12,08	11,20	12,52	12,18	13,27
Ermelo ⁽¹⁰⁾	7,98	7,90	8,72	8,54	8,58	10,34	9,93	8,11	7,47	7,68
Greytown ⁽²⁾	9,89	8,78	8,98	11,20	10,85	10,79	11,33	9,78	9,87	10,76
Greytown ⁽²⁾	8,41	7,78	8,35	8,08	9,14	9,20	9,64	8,65	8,05	9,75
Hendrina ⁽³⁾	8,78	9,64	6,63	9,07	8,54	8,49	8,89	6,73	8,35	7,13
Kinross ⁽⁸⁾	9,36	8,31	10,03	10,27	10,38	10,12	11,14	10,67	12,04	12,57
Middleburg ⁽⁵⁾	8,50	8,53	7,51	10,08	10,77	8,11	9,11	12,42	11,23	11,16
Moorivier ⁽⁷⁾	9,56	8,06	8,99	9,48	8,16	9,55	10,96	10,99	11,10	11,15
Ogies ⁽³⁾	10,92	9,67	10,33	10,46	12,29	9,45	12,59	10,18	12,84	11,39
Petit ⁽³⁾	8,49	9,59	8,89	8,54	9,58	10,04	10,69	8,67	10,38	9,29
Villiers ⁽⁸⁾	8,84	9,14	11,49	8,40	9,03	9,33	9,27	8,90	10,11	11,67
Winterton ⁽⁷⁾	13,83	12,69	13,70	13,76	14,84	14,61	15,19	13,25	15,26	13,99
Wonderfontein ⁽²⁾	8,07	6,68	6,83	6,96	5,57	7,76	9,70	7,51	6,94	8,22
2014/2015 season										
Babsfontein ⁽⁵⁾	4,61	5,09	5,51	4,92	5,43	4,33	3,98	5,10	5,68	5,33
Bergville ⁽¹⁰⁾	7,99	7,84	7,13	11,66	11,56	9,81	10,54	10,02	7,87	11,20
Bergville ⁽²⁾	12,16	11,57	12,69	12,14	13,43	12,05	13,57	12,07	14,29	15,43
Bethlehem ⁽¹⁾	3,85	5,14	4,59	5,21	5,76	5,65	4,38	5,64	6,01	5,86
Carolina ⁽⁷⁾	7,22	8,36	8,96	9,26	10,63	10,87	8,50	9,48	11,23	10,35
Delmas ⁽³⁾	8,48	8,78	9,33	8,64	8,73	8,22	7,29	9,07	10,36	7,52
Delmas ⁽²⁾	8,14	8,68	7,80	9,31	8,69	9,28	9,17	8,93	8,94	10,75
Devon ⁽⁷⁾	4,23	5,56	5,26	5,94	6,38	5,88	5,41	5,17	4,95	6,35
Greytown ⁽²⁾	5,38	5,73	5,02	6,52	3,56	5,98	5,83	4,91	4,07	5,99
Kinross ⁽⁸⁾	7,06	6,32	6,23	8,14	8,02	9,68	8,98	6,17	8,75	8,46
Middleburg ⁽⁵⁾	9,53	9,80	10,69	10,70	12,15	11,93	11,86	12,86	9,91	12,71
Moorivier ⁽⁷⁾	6,33	6,16	8,97	7,98	6,73	8,26	8,24	7,76	9,50	9,07
Standerton ⁽²⁾	3,91	3,29	3,90	3,40	3,92	3,88	4,24	3,52	3,86	3,17
Vrede ⁽⁸⁾	4,82	5,69	2,86	3,57	1,66	5,70	3,96	4,37	6,12	3,50
Winterton ⁽⁷⁾	12,21	14,25	11,98	15,86	12,84	11,93	13,45	13,14	12,56	12,42
Wonderfontein ⁽²⁾	12,61	11,23	9,11	13,15	13,21	13,89	12,98	14,18	13,42	12,83
2015/2016 season										
Babsfontein ⁽⁵⁾	5,63	6,76	5,89	6,84	5,55	7,52	5,65	6,96	6,55	7,09
Bethlehem ⁽¹⁾	4,38	6,40	4,27	5,20	5,81	5,96	4,88	4,90	5,52	5,77
Bloekomspruit ⁽¹⁾	4,28	5,87	5,47	4,88	5,51	6,40	4,97	5,57	5,99	5,96
Carolina ⁽⁷⁾	9,86	7,99	8,44	7,43	8,52	11,01	7,52	11,15	9,68	10,77
Cedara ⁽¹⁾	2,94	3,02	2,92	2,84	3,40	2,94	3,38	3,17	2,90	3,28
Delmas ⁽³⁾	7,24	7,12	8,48	7,16	8,72	8,10	8,42	10,27	8,49	8,34
Delmas ⁽⁶⁾	6,19	4,90	6,14	6,62	6,62	6,53	5,73	5,39	5,87	5,92
Dundee ⁽⁹⁾	6,96	5,90	6,78	8,38	7,70	6,59	5,78	6,89	7,25	5,34
Ermelo ⁽⁸⁾	3,19	2,88	2,63	3,56	3,12	3,71	3,10	3,62	3,74	3,34
Ermelo ⁽³⁾	1,84	3,06	3,96	2,84	4,38	2,63	2,76	5,61	2,24	3,35
Fouriesburg ⁽⁴⁾	5,04	4,97	6,86	5,18	6,06	6,58	5,32	3,97	4,83	4,07
Greytown ⁽⁸⁾	3,28	4,62	4,00	4,42	5,90	4,95	4,27	5,55	4,37	5,67
Kokstad ⁽⁹⁾	5,27	4,11	4,11	5,05	5,18	4,45	5,49	5,75	4,71	5,26
Middleburg ⁽³⁾	8,90	10,97	9,07	10,31	10,36	10,79	9,75	9,93	11,10	10,32
Petit ⁽³⁾	9,00	7,85	8,68	7,97	11,47	9,42	11,59	9,54	8,94	10,26
Wonderfontein ⁽²⁾	9,01	5,05	5,71	6,63	6,38	9,04	7,20	8,19	7,91	7,20
Gemiddelde/ Average (t ha ⁻¹)	7,57	7,56	7,62	7,99	8,29	8,38	8,32	8,27	8,39	8,47

⁽¹⁾ ARC; ⁽²⁾ Pannar; ⁽³⁾ Monsanto; ⁽⁴⁾ Pioneer; ⁽⁵⁾ Klein Karoo; ⁽⁷⁾ Seedco; ⁽⁸⁾ Link Seed; ⁽⁹⁾ DoA; ⁽¹⁰⁾ Agricol

Tabel 4 Verilog..
Table 4 Continue...

Omgeving/ Environment	Genotypes/Genotypes						Gemiddelde/ Average (t ha ⁻¹)
	PAN6Q- 408CB	PAN6R-680R	PAN6R- 880CBGT	Phb31M09	Phb33H52B	SC506	
2013/2014 season							
Babsfontein ⁽⁵⁾	10,37	11,01	9,99	10,06	9,30	9,35	4,76
Bergville ⁽²⁾	5,51	7,11	6,79	6,96	6,47	6,99	9,31
Bethlehem ⁽¹⁾	13,97	13,00	9,71	10,49	12,00	11,45	4,82
Bethlehem ⁽³⁾	5,51	6,00	5,74	8,37	6,42	4,15	7,00
Bloekomspruit ⁽¹⁾	9,60	11,22	10,42	9,50	9,30	6,33	5,55
Carolina ⁽⁷⁾	8,88	8,56	7,94	6,35	6,97	6,12	8,19
Delmas ⁽²⁾	12,24	11,29	11,09	8,10	10,92	10,32	9,10
Delmas ⁽³⁾	13,39	12,22	11,92	12,84	11,23	11,09	4,81
Devon ⁽⁸⁾	7,64	7,00	5,15	6,34	5,72	6,73	5,24
Dundee ⁽⁹⁾	7,53	6,79	5,93	5,45	5,42	7,32	8,89
Ermelo ⁽⁴⁾	13,02	12,10	11,35	10,38	9,80	9,85	6,73
Ermelo ⁽¹⁰⁾	8,47	9,67	8,76	8,81	7,43	6,87	7,92
Greytown ⁽²⁾	10,62	10,06	10,40	10,59	9,78	8,97	6,78
Greytown ⁽²⁾	9,54	9,29	9,08	8,62	9,34	8,87	6,51
Hendrina ⁽³⁾	8,50	9,37	6,44	6,34	5,90	5,19	8,12
Kinross ⁽⁸⁾	11,51	13,12	10,40	9,08	9,11	9,36	7,18
Middleburg ⁽⁵⁾	12,85	13,90	8,70	9,11	8,88	8,59	7,84
Moorivier ⁽⁷⁾	11,19	11,57	9,08	9,48	8,93	12,02	8,92
Ogies ⁽³⁾	13,54	14,03	12,48	10,70	10,63	10,20	7,48
Petit ⁽³⁾	9,53	9,46	10,02	9,01	8,85	8,64	7,58
Villiers ⁽⁸⁾	9,71	9,85	6,80	7,16	8,01	9,33	10,79
Winterton ⁽⁷⁾	13,04	14,54	12,32	12,85	14,27	13,58	5,74
Wonderfontein ⁽²⁾	9,99	8,55	6,99	8,31	6,06	7,35	0,00
2014/2015 season							
Babsfontein ⁽⁵⁾	4,61	4,36	5,89	4,92	4,63	4,96	7,02
Bergville ⁽¹⁰⁾	10,61	10,12	9,24	8,22	8,30	7,78	9,52
Bergville ⁽²⁾	13,66	12,52	11,33	13,29	13,77	10,49	4,19
Bethlehem ⁽¹⁾	6,19	5,90	5,16	5,17	4,37	4,73	7,51
Carolina ⁽⁷⁾	11,10	9,95	9,17	9,64	9,29	9,52	6,79
Delmas ⁽³⁾	10,84	8,36	10,33	8,28	7,98	8,01	7,02
Delmas ⁽²⁾	9,49	8,96	9,81	8,18	7,18	7,24	4,31
Devon ⁽⁷⁾	6,33	6,90	7,01	6,11	5,73	5,71	4,49
Greytown ⁽²⁾	6,05	5,20	5,63	5,27	4,97	5,05	5,93
Kinross ⁽⁸⁾	7,82	8,35	8,40	8,51	7,35	5,65	8,61
Middleburg ⁽⁵⁾	11,90	13,02	12,36	11,77	9,22	8,44	6,14
Moorivier ⁽⁷⁾	7,56	7,32	5,87	7,70	7,97	7,20	2,80
Standerton ⁽²⁾	3,18	3,77	3,53	4,07	3,68	2,97	3,29
Vrede ⁽⁸⁾	5,12	2,29	4,30	4,77	1,53	5,13	9,98
Winterton ⁽⁷⁾	13,29	13,48	12,50	11,82	12,64	11,66	9,57
Wonderfontein ⁽²⁾	13,99	13,61	12,07	11,95	11,24	12,20	0,00
2015/2016 season							
Babsfontein ⁽⁵⁾	7,03	5,74	6,86	6,26	6,49	5,73	4,22
Bethlehem ⁽¹⁾	5,90	5,91	5,25	4,51	3,94	4,85	4,60
Bloekomspruit ⁽¹⁾	6,28	7,42	6,68	4,66	4,27	4,22	7,18
Carolina ⁽⁷⁾	10,89	12,62	9,30	8,71	9,12	9,35	2,31
Cedara ⁽¹⁾	3,01	3,28	3,64	2,38	2,89	3,00	5,94
Delmas ⁽³⁾	10,41	8,65	8,60	7,39	9,28	5,81	4,64
Delmas ⁽⁶⁾	7,62	6,07	6,34	7,42	7,87	6,58	5,29
Dundee ⁽⁹⁾	7,55	5,54	7,59	6,18	6,44	5,60	2,57
Ermelo ⁽⁸⁾	3,48	3,31	3,55	3,47	3,04	4,08	2,68
Ermelo ⁽³⁾	3,93	3,36	5,16	2,99	4,11	2,61	4,40
Fouriesburg ⁽⁴⁾	4,61	4,59	6,54	7,05	6,21	5,16	3,89
Greytown ⁽⁸⁾	6,28	5,13	4,47	4,54	5,94	5,52	3,92
Kokstad ⁽⁹⁾	5,22	5,34	5,71	4,24	6,64	5,72	7,76
Middleburg ⁽³⁾	9,21	9,66	7,60	10,57	10,83	9,81	7,52
Petit ⁽³⁾	8,54	7,73	10,26	9,59	8,97	8,04	5,31
Wonderfontein ⁽²⁾	7,45	9,23	7,62	6,36	5,33	7,68	6,29
Gemiddelde/ Average(t ha ⁻¹)	8,82	8,68	8,10	7,83	7,67	7,44	

(1) ARC; (2) Pannar; (3) Monsanto; (4) Pioneer; (5) Klein Karoo; (7) Seedco; (8) Link Seed; (9) DoA; (10) Agricol

Tabel 5: Gemiddelde persentasie plante omval vir die 2015/2016 seisoen

Table 5: Mean percentage lodged plants for the 2015/2016 season

Genotipes Genotypes	Omval /Lodged plants				Gemiddelde Average (%)
	Bloekomspruit ⁽¹⁾	Bethlehem ⁽¹⁾	Ermelo ⁽²⁾	Greytown ⁽²⁾	
BG5285	4,42	0,00	3,64	1,95	2,50
DKC68-56R	0,92	0,00	1,23	1,32	0,87
DKC68-58BR	0,00	1,69	0,00	4,90	1,65
DKC71-42	0,00	1,79	0,00	0,65	0,61
DKC71-44B	0,00	0,98	0,61	2,84	1,11
DKC73-74BRGEN	0,00	0,98	0,00	1,35	0,58
DKC74-26R	0,00	0,00	0,00	0,63	0,16
DKC76-61B	3,85	0,00	1,22	1,28	1,59
DKC78-17B	0,00	0,00	0,00	5,48	1,37
DKC78-45BRGEN	0,91	7,50	0,63	2,00	2,76
DKC78-79BR	3,60	0,00	7,84	15,07	6,63
DKC78-87B	0,91	5,22	19,05	22,93	12,03
DKC80-40BRGEN	2,94	2,68	3,67	0,00	2,32
IMP52-11R	2,65	2,54	1,74	0,61	1,89
IMP52-12R	0,00	0,00	1,85	2,67	1,13
IMP53-49B	0,91	0,00	0,00	0,69	0,40
KKS8301B	1,98	4,63	0,00	2,10	2,18
KKS8326B	0,00	5,45	5,13	0,00	2,65
KKS8408R	0,97	0,88	0,00	0,00	0,46
KKS8410BR	2,88	0,00	0,00	0,68	0,89
LS8518	8,08	14,41	0,00	2,84	6,33
LS8526	0,00	0,00	1,89	2,13	1,01
LS8533R	1,85	0,00	4,04	1,39	1,82
LS8536B	1,89	4,72	2,56	3,45	3,16
LS8537	1,87	13,33	0,00	2,58	4,45
LS8539B	8,00	5,13	0,00	1,36	3,62
LS8541BR	2,91	0,00	2,67	0,70	1,57
P2137B	2,00	0,00	1,94	7,38	2,83
P2369WB	0,94	0,00	1,26	0,65	0,71
P2432R	1,00	0,00	0,00	0,00	0,25
P2653WB	3,74	0,00	0,00	1,97	1,43
PAN4A-111	0,90	1,67	0,88	0,71	1,04
PAN4A-172	0,00	3,42	0,62	4,73	2,19
PAN4B-311B	1,96	1,68	1,76	2,56	1,99
PAN4R-511R	0,93	0,00	4,82	0,00	1,44
PAN5A-182	0,00	2,54	0,00	0,66	0,80
PAN5A-291	0,98	0,00	0,66	1,41	0,76
PAN5R-791BR	1,75	3,54	2,00	1,38	2,17
PAN6P-110	0,00	5,71	1,00	0,00	1,68
PAN6Q-245	1,82	0,00	0,00	4,64	1,62
PAN6Q-345CB	0,00	0,00	0,00	4,17	1,04
PAN6Q-408CB	3,48	0,00	1,18	2,61	1,82
PAN6R-680R	0,00	3,74	0,00	0,65	1,10
PAN6R-880CBGT	0,00	1,94	0,00	2,01	0,99
Phb31M09	1,85	3,39	0,60	0,00	1,46
Phb33H52B	0,97	1,94	1,24	0,00	1,04
SC402	5,88	5,31	4,90	6,08	5,54
SC419	9,71	4,55	9,26	3,36	6,72
SC506	3,13	5,61	3,40	3,47	3,90
SC512	0,97	0,00	0,00	2,72	0,92
Gemiddelde/ Average (%)	1,87	2,34	1,87	2,66	2,19

⁽¹⁾ ARC; ⁽²⁾ Pannar

Tabel 6: Gemiddelde persentasie spruitvorming vir die 2015/2016 seisoen**Table 6: Mean percentage tillering for the 2015/2016 season**

Genotipes Genotypes	Spruite /Tillering (%)					Gemiddelde Average (%)
	Bloekomspruit ⁽¹⁾	Bethelhem ⁽¹⁾	Ermelo ⁽⁸⁾	Greytown ⁽⁸⁾	Kokstad ⁽⁹⁾	
BG5285	10,62	19,09	0,00	1,30	1,75	6,55
DKC68-56R	18,35	0,00	0,00	4,64	2,63	5,12
DKC68-58BR	15,18	6,78	2,06	8,39	0,94	6,67
DKC71-42	14,04	0,00	0,00	2,58	1,79	3,68
DKC71-44B	12,04	0,00	1,21	1,42	0,00	2,93
DKC73-74BRGEN	18,87	0,00	0,00	1,35	2,59	4,56
DKC74-26R	13,39	7,56	3,59	5,70	0,88	6,22
DKC76-61B	25,00	5,61	2,44	0,64	5,98	7,93
DKC78-17B	16,51	7,69	0,75	2,74	7,63	7,06
DKC78-45BRGEN	21,82	5,83	0,00	5,33	3,45	7,29
DKC78-79BR	17,12	2,61	1,96	6,85	0,85	5,88
DKC78-87B	13,64	6,96	1,90	8,92	0,00	6,28
DKC80-40BRGEN	14,71	0,00	1,83	4,67	1,77	4,60
IMP52-11R	23,01	8,47	5,23	14,02	5,98	11,34
IMP52-12R	14,14	13,56	0,62	3,33	6,25	7,58
IMP53-49B	25,45	22,50	5,83	24,14	4,24	16,43
KKS8301B	16,83	7,41	0,00	0,00	2,70	5,39
KKS8326B	17,00	0,00	5,77	5,59	6,90	7,05
KKS8408R	18,45	8,85	0,00	2,16	0,00	5,89
KKS8410BR	8,65	0,00	0,00	1,35	5,31	3,06
LS8518	13,13	0,00	0,00	2,84	0,00	3,19
LS8526	19,80	5,13	0,00	13,48	1,77	8,04
LS8533R	15,74	8,55	1,01	4,17	8,85	7,66
LS8536B	19,81	9,43	2,56	5,52	0,88	7,64
LS8537	23,36	6,67	1,23	2,58	1,68	7,11
LS8539B	15,00	0,00	0,65	1,36	0,00	3,40
LS8541BR	18,45	0,00	0,67	7,04	3,54	5,94
P2137B	23,00	13,33	0,65	3,36	1,75	8,42
P2369WB	21,70	2,59	3,77	1,94	0,00	6,00
P2432R	10,00	18,26	2,52	1,21	6,84	7,77
P2653WB	27,10	29,17	0,00	9,21	11,21	15,34
PAN4A-111	19,82	10,00	1,77	10,64	5,88	9,62
PAN4A-172	14,29	2,56	0,00	5,41	1,67	4,78
PAN4B-311B	20,59	9,24	1,18	10,26	5,98	9,45
PAN4R-511R	20,37	26,67	0,00	11,54	5,08	12,73
PAN5A-182	11,97	5,93	5,94	3,97	0,88	5,74
PAN5A-291	13,73	16,10	0,66	4,23	0,00	6,94
PAN5R-791BR	11,40	0,88	0,00	4,14	0,00	3,28
PAN6P-110	21,78	0,00	0,00	1,90	0,00	4,74
PAN6Q-245	10,91	13,33	1,85	1,99	1,71	5,96
PAN6Q-345CB	21,43	9,09	1,20	4,17	3,51	7,88
PAN6Q-408CB	20,87	6,42	0,00	0,65	0,84	5,76
PAN6R-680R	18,35	8,41	5,17	6,54	0,86	7,87
PAN6R-880CBGT	20,62	15,53	6,41	0,00	0,90	8,69
Phb31M09	13,89	0,00	0,60	7,84	0,00	4,47
Phb33H52B	19,42	5,83	1,24	0,00	0,00	5,30
SC402	17,65	0,00	0,00	2,03	0,00	3,94
SC419	13,59	2,73	0,00	5,04	1,69	4,61
SC506	16,67	2,80	0,00	6,25	3,45	5,83
SC512	15,53	5,22	0,65	0,68	0,84	4,58
Gemiddelde/ Average (%)	17,30	7,14	1,46	4,90	2,63	6,68

⁽¹⁾ ARC; ⁽⁸⁾ Link Seed; ⁽⁹⁾ DoA

Tabel 7: Gemiddelde aantal koppe per plant vir die 2015/2016 seisoen

Table 7: Mean number of ears per plant for the 2015/2016 season

Genotypes Genotypes	Koppe / Ears plant ⁻¹				Gemiddelde Average (plant ⁻¹)
	Bloekomspruit ⁽¹⁾	Bethelhem ⁽¹⁾	Dundee ⁽⁹⁾	Kokstad ⁽⁹⁾	
BG5285	1,56	1,87	1,05	1,18	1,42
DKC68-56R	1,37	1,34	1,10	1,00	1,20
DKC68-58BR	1,28	1,47	0,94	1,11	1,20
DKC71-42	1,22	1,26	1,26	1,01	1,19
DKC71-44B	1,25	1,60	1,03	1,01	1,22
DKC73-74BRGEN	1,25	0,66	0,85	0,96	0,93
DKC74-26R	1,34	1,58	1,09	1,04	1,26
DKC76-61B	1,53	1,45	0,64	1,07	1,17
DKC78-17B	1,73	1,62	1,07	1,45	1,47
DKC78-45BRGEN	1,95	1,64	0,97	1,58	1,53
DKC78-79BR	1,76	1,54	1,15	1,41	1,47
DKC78-87B	1,61	1,70	1,26	1,43	1,50
DKC80-40BRGEN	1,91	1,56	1,22	1,24	1,48
IMP52-11R	1,27	1,11	0,96	0,97	1,08
IMP52-12R	1,31	1,00	0,66	0,96	0,98
IMP53-49B	1,35	1,36	1,02	1,04	1,19
KKS8301B	1,37	0,78	0,68	1,02	0,96
KKS8326B	1,22	1,12	1,01	1,01	1,09
KKS8408R	1,40	1,16	1,10	0,97	1,16
KKS8410BR	1,18	1,08	1,17	1,00	1,11
LS8518	1,30	0,91	1,09	1,00	1,07
LS8526	1,20	0,89	0,92	0,84	0,96
LS8533R	1,57	1,62	1,11	1,06	1,34
LS8536B	1,36	1,13	0,99	0,95	1,11
LS8537	1,31	1,23	1,12	0,98	1,16
LS8539B	1,44	1,30	1,13	1,12	1,25
LS8541BR	1,64	1,67	1,21	1,00	1,38
P2137B	1,40	1,29	1,19	1,09	1,24
P2369WB	1,42	1,12	0,48	0,96	0,99
P2432R	1,19	1,14	1,01	0,98	1,08
P2653WB	1,31	1,15	0,95	1,08	1,12
PAN4A-111	1,11	1,21	1,17	0,99	1,12
PAN4A-172	1,24	0,62	0,35	0,98	0,80
PAN4B-311B	1,25	1,07	0,96	0,99	1,07
PAN4R-511R	1,31	1,06	0,97	1,08	1,11
PAN5A-182	1,64	1,70	1,38	1,28	1,50
PAN5A-291	1,53	1,46	1,13	1,03	1,29
PAN5R-791BR	1,71	1,64	1,61	1,08	1,51
PAN6P-110	1,59	1,48	1,13	1,40	1,40
PAN6Q-245	1,65	1,25	1,07	1,40	1,34
PAN6Q-345CB	1,47	1,74	0,89	1,47	1,39
PAN6Q-408CB	1,61	1,57	1,32	1,15	1,41
PAN6R-680R	1,76	1,64	1,23	1,40	1,51
PAN6R-880CBGT	1,58	1,11	1,14	1,22	1,26
Phb31M09	1,14	1,00	0,94	0,89	0,99
Phb33H52B	1,19	1,22	1,03	0,96	1,10
SC402	1,21	1,16	1,09	1,24	1,18
SC419	1,30	1,20	1,13	1,08	1,18
SC506	1,18	1,13	1,06	0,97	1,09
SC512	1,17	1,10	0,72	0,98	0,99
Gemiddelde/ Average (plant ⁻¹)	1,41	1,29	1,04	1,10	1,21

⁽¹⁾ ARC; ⁽⁹⁾ DoA

Tabel 8: Gemiddelde graanvogpersentasie met oestyd 2015/2016 seisoen

Table 8: Mean percentage grain moisture at harvest for the 2015/2016 season

Genotipes Genotypes	Graan vog /Grain moisture (%)									Gemiddelde/ Average (%)
	Villers ⁽⁵⁾	Ermelo ⁽³⁾	Dundee ⁽⁹⁾	Bethelhem ⁽¹⁾	Ermelo ⁽⁸⁾	Delmas ⁽³⁾	Peti ⁽³⁾	Kokstad ⁽⁹⁾	Kinross ⁽⁸⁾	
BG5285	12,8	19,0	12,9	16,0	28,6	12,0	23,5	15,8	26,1	18,52
DKC68-56R	13,4	21,8	12,8	14,4	22,8	10,5	18,8	11,8	19,4	16,19
DKC68-58BR	13,1	18,4	13,3	15,0	27,4	11,8	18,8	12,1	20,2	16,68
DKC71-42	12,8	19,4	13,2	11,7	19,9	10,3	15,5	9,5	16,3	14,29
DKC71-44B	12,6	17,1	12,4	11,8	19,8	10,7	16,6	10,3	14,0	13,92
DKC73-74BRGEN	15,7	25,1	13,0	18,6	25,0	13,4	19,9	16,3	23,4	18,93
DKC74-26R	13,8	23,1	12,6	15,3	27,5	13,3	20,0	15,8	23,0	18,27
DKC76-61B	15,1	25,1	12,0	18,5	24,5	12,7	24,9	16,1	28,3	19,69
DKC78-17B	14,8	25,2	13,5	17,4	26,6	12,7	24,0	16,1	26,3	19,62
DKC78-45BRGEN	15,0	21,5	13,2	21,0	27,4	13,4	24,6	15,5	24,8	19,60
DKC78-79BR	14,6	20,6	12,2	17,3	25,1	12,9	22,1	16,7	23,9	18,38
DKC78-87B	15,1	20,2	13,2	15,4	24,8	11,6	20,0	14,7	26,8	17,98
DKC80-40BRGEN	13,0	22,0	12,8	15,1	24,3	12,8	22,7	14,5	22,9	17,79
IMP52-11R	12,4	20,9	12,4	15,2	22,6	11,5	17,4	13,7	18,2	16,03
IMP52-12R	12,9	22,0	12,4	16,7	20,2	11,4	17,6	12,4	15,7	15,70
IMP53-49B	13,8	23,7	13,4	17,5	26,5	12,9	24,0	17,8	24,1	19,30
KKS8301B	13,0	23,2	11,7	12,9	21,3	11,3	17,5	9,8	22,2	15,88
KKS8326B	14,2	19,6	14,1	15,7	26,9	13,8	18,2	15,8	25,6	18,21
KKS8408R	11,3	18,8	12,5	17,1	25,8	12,2	18,9	15,3	22,6	17,17
KKS8410BR	13,2	20,6	13,2	17,5	24,5	13,8	19,2	15,5	23,3	17,87
LS8518	13,0	20,6	12,6	12,6	23,0	11,2	15,3	12,9	14,7	15,10
LS8526	13,3	22,0	14,1	19,3	26,8	15,3	22,3	16,7	21,0	18,98
LS8533R	13,3	24,5	12,9	17,3	27,0	10,5	21,5	14,2	22,9	18,23
LS8536B	13,1	14,4	12,5	13,2	21,7	11,6	14,8	13,0	16,2	14,50
LS8537	16,6	24,2	13,4	23,1	29,4	14,0	25,5	17,2	26,0	21,04
LS8539B	12,3	24,1	12,6	17,1	28,6	11,6	24,4	16,1	24,6	19,04
LS8541BR	15,8	21,6	12,7	18,4	25,1	14,7	23,1	15,6	27,4	19,38
P2137B	13,2	19,9	12,4	14,5	20,0	11,0	16,4	10,2	14,7	14,70
P2369WB	14,7	21,8	12,2	13,8	24,2	10,5	20,0	11,5	22,0	16,74
P2432R	14,4	24,5	12,8	15,8	26,6	12,8	22,2	15,1	23,0	18,58
P2653WB	17,0	21,9	13,0	17,9	27,4	15,3	21,3	17,3	23,8	19,43
PAN4A-111	13,8	20,2	12,9	17,1	23,5	13,2	18,1	14,7	17,2	16,74
PAN4A-172	13,3	21,1	12,2	13,8	20,0	11,3	17,9	12,9	16,3	15,42
PAN4B-311B	13,5	24,0	13,6	18,0	23,7	12,9	19,5	15,8	20,5	17,94
PAN4R-511R	13,8	24,2	12,5	16,2	24,3	12,8	19,3	15,6	20,3	17,67
PAN5A-182	14,4	21,9	12,6	16,3	25,1	12,5	20,4	16,0	20,1	17,70
PAN5A-291	15,2	24,5	13,8	17,4	31,4	12,4	26,8	17,4	26,8	20,63
PAN5R-791BR	15,1	23,3	12,8	19,3	29,5	12,3	26,0	16,5	26,2	20,11
PAN6P-110	15,7	25,6	12,9	17,6	22,2	13,9	23,4	16,9	26,1	19,37
PAN6Q-245	14,6	25,9	13,0	18,5	26,4	11,7	22,7	17,3	28,8	19,88
PAN6Q-345CB	13,5	25,8	13,2	16,7	27,2	13,4	20,8	16,0	28,6	19,47
PAN6Q-408CB	16,5	23,5	14,6	19,2	23,5	12,3	23,2	16,2	28,6	19,73
PAN6R-680R	18,0	24,5	14,2	20,0	27,2	13,7	25,6	17,6	24,1	20,54
PAN6R-880CBGT	13,5	23,0	13,4	16,5	28,3	14,6	22,0	16,4	23,4	19,01
Phb31M09	12,3	15,8	12,3	16,8	20,6	11,3	17,5	11,9	19,3	15,31
Phb33H52B	13,3	19,4	13,1	14,2	16,5	10,5	15,3	9,6	14,6	14,06
SC402	14,3	20,5	13,4	17,2	22,1	14,3	21,9	15,2	19,8	17,63
SC419	14,3	24,2	13,6	20,1	21,9	15,1	22,9	17,6	21,6	19,03
SC506	17,1	22,6	13,6	16,1	23,2	14,3	20,9	16,3	26,0	18,90
SC512	13,2	23,7	12,8	14,0	20,9	13,7	16,9	11,9	17,3	16,04
Gemiddelde/ Average (%)	14,1	22,0	13,0	16,6	24,6	12,6	20,6	14,7	22,2	

⁽¹⁾ ARC; ⁽³⁾ Monsanto; ⁽⁵⁾ Klein Karoo; ⁽⁸⁾ Link Seed; ⁽⁹⁾ DoA

Tabel 9: Gemiddelde waardes vir alle agronomiese parameters vir die 2015/2016 seisoen

Table 9: Summary of mean values for all agronomic parameters for the 2015/2016 season

Genotipes Genotypes	Omval % Lodged %			Spruite % Tillering %	Koppe (plant ⁻¹) Ears (plant ⁻¹)	Graan vog % Grain moisture %	Graan obrengs Grain yield (t ha ⁻¹)
	*	Stalk	Root				
BG5285	6,92	2,97	0,00	6,55	1,42	17,3	6,32
DKC68-56R	9,83	0,77	0,00	5,12	1,20	15,4	6,60
DKC68-58BR	9,63	4,13	0,17	6,67	1,20	15,9	6,44
DKC71-42	4,99	0,30	0,17	3,68	1,19	13,8	6,50
DKC71-44B	8,41	1,27	0,00	2,93	1,22	13,5	6,61
DKC73-74BRGEN	6,50	0,97	0,47	4,56	0,93	17,7	6,32
DKC74-26R	2,25	0,64	0,00	6,22	1,26	17,1	6,72
DKC76-61B	10,52	0,93	0,65	7,93	1,17	18,3	6,24
DKC78-17B	2,96	3,87	0,33	7,06	1,47	18,2	6,50
DKC78-45BRGEN	6,66	3,09	0,00	7,29	1,53	18,2	6,45
DKC78-79BR	9,49	2,80	0,00	5,88	1,47	17,2	6,68
DKC78-87B	13,93	3,15	0,00	6,28	1,50	16,9	6,73
DKC80-40BRGEN	5,29	1,13	0,53	4,60	1,48	16,7	6,79
IMP52-11R	2,97	0,77	0,00	11,34	1,08	15,2	6,04
IMP52-12R	8,39	0,80	0,20	7,58	0,98	14,9	5,74
IMP53-49B	3,75	1,23	0,00	16,43	1,19	18,0	6,36
KKS8301B	7,13	0,37	0,20	5,39	0,96	15,2	5,73
KKS8326B	3,31	1,77	0,60	7,05	1,09	17,1	6,01
KKS8408R	5,39	0,80	0,00	5,89	1,16	16,1	6,26
KKS8410BR	2,04	3,23	0,00	3,06	1,11	16,8	6,58
LS8518	17,66	4,13	0,35	3,19	1,07	14,4	5,78
LS8526	5,01	1,57	0,00	8,04	0,96	17,6	5,82
LS8533R	4,43	1,84	0,20	7,66	1,34	17,3	5,72
LS8536B	14,51	2,83	0,00	7,64	1,11	14,0	5,86
LS8537	6,57	4,17	0,73	7,11	1,16	19,5	6,05
LS8539B	7,45	1,83	0,57	3,40	1,25	17,7	5,95
LS8541BR	4,26	2,65	0,00	5,94	1,38	18,0	5,79
P2137B	6,56	1,33	0,00	8,42	1,24	14,2	6,85
P2369WB	7,36	0,30	0,64	6,00	0,99	16,0	6,33
P2432R	2,85	0,61	0,15	7,77	1,08	17,4	6,54
P2653WB	5,57	2,55	0,00	15,34	1,12	18,1	6,67
PAN4A-111	3,40	2,79	0,00	9,62	1,12	15,8	6,06
PAN4A-172	5,65	1,60	0,00	4,78	0,80	14,7	6,33
PAN4B-311B	4,94	1,83	0,00	9,45	1,07	16,7	5,99
PAN4R-511R	3,02	2,77	0,00	12,73	1,11	16,6	6,53
PAN5A-182	2,96	0,33	0,00	5,74	1,50	16,6	7,01
PAN5A-291	4,50	1,07	0,47	6,94	1,29	19,0	6,46
PAN5R-791BR	6,18	2,40	0,50	3,28	1,51	18,5	6,51
PAN6P-110	5,25	2,80	0,16	4,74	1,40	18,1	6,73
PAN6Q-245	2,61	6,10	0,00	5,96	1,34	18,5	6,26
PAN6Q-345CB	7,02	3,73	0,00	7,88	1,39	18,1	6,38
PAN6Q-408CB	3,95	0,67	0,16	5,76	1,41	18,4	6,72
PAN6R-680R	2,91	0,97	0,00	7,87	1,51	19,1	6,38
PAN6R-880CBGT	2,44	0,00	0,00	8,69	1,26	17,9	6,66
Phb31M09	6,49	1,72	0,00	4,47	0,99	14,6	6,02
Phb33H52B	2,55	0,00	0,30	5,30	1,10	13,6	6,40
SC402	11,24	2,00	0,34	3,94	1,18	16,6	5,27
SC419	8,44	3,82	1,14	4,61	1,18	17,7	5,59
SC506	8,00	1,65	1,07	5,83	1,09	17,7	5,85
SC512	8,95	1,78	0,00	4,58	0,99	15,3	5,47
Gemiddelde/Average	6,26	1,94	0,20	6,68	1,21	16,7	

*unspecified type of lodging

Tabel 10: Meerjarige gemiddeldes ten opsigte van persentasie plante omval vir die 2013/2014, 2014/2015 & 2015/2016 seisoene

Table 10: Multi seasonal means for total percentage lodged plants for the 2013/2014, 2014/2015 & 2015/2016 seasons

Genotipes Genotypes	Omval / Lodged (%)			Gemiddelde/ Average (%)
	2013/2014	2014/2015	2015/2016	
BG5285	9,62	2,00	2,72	4,78
DKC73-74BRGEN	5,81	1,80	0,91	2,84
DKC78-17B	4,95	3,40	2,57	3,64
DKC78-45BRGEN	4,89	1,70	2,86	3,15
DKC78-79BR	7,25	3,60	4,99	5,28
DKC78-87B	4,81	3,50	11,94	6,75
DKC80-40BRGEN	10,87	4,30	1,96	5,71
IMP52-11R	4,81	5,30	1,41	3,84
KKS8410BR	5,81	4,90	1,92	4,21
LS8518	4,35	2,80	5,48	4,21
LS8526	13,36	5,10	1,25	6,57
LS8533R	5,49	4,00	1,85	3,78
LS8536B	10,51	5,50	3,04	6,35
LS8539B	10,64	1,80	2,89	5,11
P2432R	3,87	1,90	0,47	2,08
P2653WB	5,78	4,90	1,92	4,20
PAN4B-311B	5,97	2,20	5,78	4,65
PAN6P-110	5,85	3,50	2,26	3,87
PAN6Q-245	9,94	4,60	3,52	6,02
PAN6Q-345CB	8,48	4,70	2,18	5,12
PAN6Q-408CB	6,45	3,50	1,33	3,76
PAN6R-680R	7,88	2,70	1,03	3,87
PAN6R-880CBGT	5,07	1,10	0,55	2,24
Phb31M09	7,37	4,20	1,51	4,36
Phb33H52B	4,44	1,70	0,55	2,23
SC506	5,45	3,90	3,40	4,25
Gemiddelde/Average (%)	6,91	3,41	2,70	

Tabel 11: Meerjarige gemiddeldes ten opsigte van persentasie spruitvorming vir die 2013/2014, 2014/2015 & 2015/2016 seisoene

Table 11: Multi seasonal means for total percentage tillering for the 2013/2014, 2014/2015 & 2015/2016 seasons

Genotipes Genotypes	Spruite/ Tillering (%)			Gemiddelde Average (%)
	2013/2014	2014/2015	2015/2016	
BG5285	4,00	8,40	6,55	6,30
DKC73-74BRGEN	3,88	4,50	4,56	4,32
DKC78-17B	7,67	11,20	7,06	8,64
DKC78-45BRGEN	7,43	10,60	7,29	8,42
DKC78-79BR	5,06	8,70	5,88	6,55
DKC78-87B	4,97	9,90	6,28	7,05
DKC80-40BRGEN	5,76	6,80	4,60	5,71
IMP52-11R	7,40	15,30	11,34	11,35
KKS8410BR	3,63	6,60	3,06	4,44
LS8518	4,77	9,10	3,19	5,68
LS8526	2,99	5,30	8,04	5,44
LS8533R	4,12	10,80	7,66	7,52
LS8536B	3,22	5,60	7,64	5,48
LS8539B	4,84	8,00	3,40	5,41
P2432R	6,16	11,60	7,77	8,51
P2653WB	7,62	14,80	15,34	12,58
PAN4B-311B	7,03	14,70	9,45	9,74
PAN6P-110	4,95	11,80	4,74	7,16
PAN6Q-245	4,18	11,30	5,96	7,15
PAN6Q-345CB	6,47	11,40	7,88	8,57
PAN6Q-408CB	5,02	11,50	5,76	7,43
PAN6R-680R	6,12	10,90	7,87	8,30
PAN6R-880CBGT	5,50	12,70	8,69	8,96
Phb31M09	5,08	11,70	4,47	7,08
Phb33H52B	5,53	11,10	5,3	7,31
SC506	4,74	6,00	5,83	5,51
Gemiddelde/Average (%)	5,13	10,01	6,75	

Tabel 12 Meerjarige gemiddeldes ten opsigte van aantal koppe per plant vir die 2013/2014, 2014/2015 & 2015/2016 seisoene

Table 12: Multi seasonal means for number of ears per plant for the 2013/2014, 2014/2015 & 2015/2016 seasons

Genotipes Genotypes	Koppe / Ears (plant ⁻¹)			Gemiddelde/ Average (%)
	2013/2014	2014/2015	2015/2016	
BG5285	1,40	1,26	1,42	1,36
DKC73-74BRGEN	1,13	1,06	0,93	1,04
DKC78-17B	1,54	1,38	1,47	1,46
DKC78-45BRGEN	1,49	1,56	1,53	1,53
DKC78-79BR	1,52	1,44	1,47	1,48
DKC78-87B	1,18	1,44	1,50	1,37
DKC80-40BRGEN	1,66	1,37	1,48	1,50
IMP52-11R	1,18	1,27	1,08	1,18
KKS8410BR	1,27	1,08	1,11	1,15
LS8518	1,20	1,12	1,07	1,13
LS8526	1,58	1,42	0,96	1,32
LS8533R	1,45	1,28	1,34	1,36
LS8536B	1,34	1,27	1,11	1,24
LS8539B	1,21	1,18	1,25	1,21
P2432R	1,41	1,17	1,08	1,22
P2653WB	1,18	1,17	1,12	1,16
PAN4B-311B	1,16	1,01	1,07	1,08
PAN6P-110	1,57	1,38	1,4	1,45
PAN6Q-245	1,60	1,40	1,34	1,45
PAN6Q-345CB	1,60	1,37	1,39	1,45
PAN6Q-408CB	1,51	1,37	1,41	1,43
PAN6R-680R	1,67	1,44	1,51	1,54
PAN6R-880CBGT	1,52	1,26	1,26	1,35
Phb31M09	1,19	1,08	0,99	1,09
Phb33H52B	1,36	1,19	1,1	1,22
SC506	1,24	1,05	1,09	1,13
Gemiddelde/Average (plant ⁻¹)	1,39	1,27	1,25	

Tabel 13: Meerjarige gemiddeldes ten opsigte van persentasie graanvog vir die 2013/2014, 2014/2015 & 2015/2016 seisoene

Table 13: Multi seasonal means for percentage grain moisture for the 2013/2014, 2014/2015 & 2015/2016 seasons

Genotipes Genotypes	Graan vog/ Grain moisture (%)			Gemiddelde Average (%)
	2013/2014	2014/2015	2015/2016	
BG5285	15,03	16,50	18,5	16,68
DKC73-74BRGEN	15,14	16,10	18,9	16,71
DKC78-17B	15,40	16,70	19,6	17,23
DKC78-45BRGEN	15,21	16,70	19,6	17,17
DKC78-79BR	14,81	15,00	18,4	16,07
DKC78-87B	15,08	16,40	18,0	16,49
DKC80-40BRGEN	14,87	16,30	17,8	16,32
IMP52-11R	14,27	15,20	16,0	15,16
KKS8410BR	14,07	16,10	17,9	16,02
LS8518	15,14	17,10	15,1	15,78
LS8526	12,42	13,50	19,0	14,97
LS8533R	13,79	15,90	18,2	15,96
LS8536B	13,76	14,50	14,5	14,25
LS8539B	15,35	16,90	19,0	17,08
P2432R	14,64	16,20	18,6	16,48
P2653WB	15,96	16,80	19,4	17,39
PAN4B-311B	15,02	16,30	17,9	16,41
PAN6P-110	16,16	17,80	19,4	17,79
PAN6Q-245	15,92	176,20	19,9	70,67
PAN6Q-345CB	15,24	16,20	19,5	16,98
PAN6Q-408CB	15,72	17,80	19,7	17,74
PAN6R-680R	17,36	18,50	20,5	18,79
PAN6R-880CBGT	15,47	16,20	19,0	16,89
Phb31M09	13,25	14,40	15,3	14,32
Phb33H52B	13,02	13,90	14,1	13,67
SC506	15,97	16,80	18,9	17,22
Gemiddelde/Average (%)	14,93	22,31	18,18	

Tabel 14: Opsomming van gemiddelde waardes vir alle agronomiese parameters vir die 2013/2014, 2014/2015 & 2015/2016 seisoene

Table 14: Summary of mean values for all agronomic parameters for the 2013/2014, 2014/2015 & 2015/2016 seasons

Genotipes Genotypes	Omvul % Logged %	Spruite % Tillering %	Koppe plant ⁻¹ Ears plant ⁻¹	Graan vog% Grain moisture%	Graan opbrengs Grain yield (t ha ⁻¹)
BG 5285	4,78	6,30	1,36	16,7	8,35
DKC 73 - 74BR GEN	2,84	4,32	1,04	16,7	8,32
DKC 78 - 17B	3,64	8,64	1,46	17,3	8,42
DKC 78 - 45BR GEN	3,15	8,42	1,53	17,3	7,96
DKC 78 - 79BR	5,28	6,55	1,48	16,6	8,44
DKC 78 - 87B	6,75	7,05	1,52	16,5	8,38
DKC 80 - 40BR GEN	5,71	5,71	1,50	16,3	8,54
IMP 52 - 11R	3,84	11,35	1,13	15,2	8,05
KKS 8410BR	4,21	4,44	1,15	16,0	8,04
LS 8518	4,21	5,68	1,13	15,8	8,05
LS 8526	6,57	5,44	1,32	15,0	7,57
LS 8533R	3,78	7,52	1,36	16,0	7,56
LS 8536B	6,35	5,48	1,24	14,3	7,62
LS 8539B	5,11	5,41	1,21	17,1	7,99
P 2432R	2,08	8,51	1,22	16,5	8,29
P 2653WB	4,20	12,58	1,16	17,4	8,38
PAN 4B - 311B	4,65	9,74	1,15	15,9	8,32
PAN 6P - 110	3,87	7,16	1,45	17,8	8,27
PAN 6Q - 245	6,02	7,15	1,45	17,7	8,39
PAN 6Q - 345CB	5,12	8,57	1,45	17,0	8,47
PAN 6Q - 408CB	3,76	7,43	1,43	17,7	8,82
PAN 6R -680R	3,87	8,30	1,54	18,8	8,68
PAN 6R - 880CB GT	2,24	8,96	1,35	16,9	8,10
Phb 31M09	4,36	7,08	1,09	14,3	7,83
Phb 33H52B	2,23	7,31	1,22	13,7	7,67
SC 506	4,25	5,51	1,13	17,2	7,44
Gemiddelde/Average	4,34	7,33	1,31	16,4	